
8 NCBI

2xh41+1\$ 09/843159

CDART: Conserved Domain Architecture Retrieval Tool



Result page: Previous

Next

Subset by Taxonomy

Subset by selected domains: pfam00644 Poly(ADP-ribose) polymerase catalytic domain. Pol... COG3831 Uncharacterized conserved protein [Function unkno... cd00204 ankyrin repeats; ankyrin repeats mediate protein... includes: COG0666 COG3779 pfam00533 BRCA1 C Terminus (BRCT) domain. The BRCT domain i... smart00292 cd00027 includes: SAP domain. The SAP (after SAF-A/B, Acinus and Pl... pfam02037 includes: smart00513 pfam02877 Poly(ADP-ribose) polymerase, regulatory domain. P... Poly(ADP-ribose) polymerase and DNA-Ligase Zn-fin... pfam00645

> Help | Disclaimer | Write to the Help Desk NCBI | NLM | NIH

Help?



Exhibit J 09/843,159

Conserved Domain Database



Structure CDD Taxonomy Protein Source: Pfam[US], Pfam[UK] CD: pfam02877.8, PARP_reg PSSM-Id: 3371 Description: Poly(ADP-ribose) polymerase, regulatory domain. Poly(ADP-ribose) polymerase catalyses the covalent attachment of ADP-ribose units from NAD+ to itself and to a limited number of other DNA binding proteins, which decreases their affinity for DNA. Poly(ADP-ribose) polymerase is a regulatory component induced by DNA damage. The carboxyl-terminal region is the most highly conserved region of the protein. Experiments have shown that a carboxyl 40 kDa fragment is still catalytically active. Taxa: Eukaryota References: 3 Pubmed Links Created: 11-Apr-2003 Status: Alignment from source PSSM: 134 columns Representative: Consensus Aligned: 6 rows Proteins: [Click here for CDART summary of Proteins containing pfam02877] View 3D Structure with Cn3D Virtual Bonds ▼ (To display structure, download Cn3D) color at 2.0 bits ▼ width 60 View Alignment as Hypertext up to 10 Subset Rows of the most diverse members 😴 30 10 20 40 50 *....|....*....|....*....|..... 1 KSKLLKSVQDLIKLIFDVDSMAQTMMEFEI--DMEKMPLGKLSKRQIQSAYRVLKEIYEV 58 consensus 9 KSKLAKPIQDLIKMIFDVESMKKAMVEFEI--DLQKMPLGKLSKRQIQSAYSILNEVQQA 66 gi 1353140 171 LLKQLK-FNEAFGRPIDCLSLAQLTTGYEIlsKIEESIGGKSARRSTRGRPRVADRVLAV 229 gi 1709740 286 QSKLDTRVAKFISLICNVSMMAQHMMEIGY--NANKLPLGKISKSTISKGYEVLKRISEV 343 644 TSKLEISVONLIKLIFDIDSMNKTLMEFHI--DMDKMPLGKLSAHQIQSAYRVVKEIYNV gi 1709741 647 KSKLPLSVQDIIKLMFDVDSMKRTMMEFDL--DMEKMFLGKLSQKQIQSAYKVLTEIYEL 704 80 90 100 110|....*.... 59 ISDGGSPAKLIDLSNRFYTLIPHDFGFKKPP--LIDTHQKIQAKRQMLDALK-EIEVAYS 115 consensus 67 VSDGGSESQILDLSNRFYTLIPHDFGMKKPF--LLSNLEYIQAKVQMLDNLL-DIEVAYS 123 3PAX 1353140 230 KSDGPS---LHDI-NKYYSLIPHSFGFCVPP--KIDSHAKIQAERELLDALKgSIEASLE 283 gi 1709740 344 I-DRYDRTRLEELSGEFYTVIPHDFGFKKMSqfVIDTPQKLKQKIEMVEALG-EIELATK 401 gi 548585 702 LECGSNTAKLIDATNRFYTLIPHNFGVQLPT--LIETHQQIEDDRQMLDSLA-EIEVAYS 758 gi 1709741 705 IQGGGTNAKFIDATNRFYTLIPHNFGTQSPP--LLDTTEQVEQLRQMLDSLI-EIECAYS 761 130 ...*...|....*... 116 LLDLEDTASDKDPLDRHYE 134 consensus 124 LLEGGNEDGDKDPIDINYE 142 qi 1353140 284 LEDLKETASSEDIYQELYE 300 gi 1709740 402 LLSVDPGLQD-DPLYYHYQ 419 gi 548585 759 IIKSEDVSDACNPLDNHYA 777 1709741 762 LLQTEDSKADINPIDKHYE 780

Help | Disclaimer | Write to the Help Desk NCB| | NLM | NIH

Help?

5xhusir J 09/843,159 2/2



Conserved Domain Database

cnn Structure Taxonomy Protein Nucleotide Source: Pfam[US], Pfam[UK] CD: pfam00644.8, PARP PSSM-Id: 1202 Description: Poly(ADP-ribose) polymerase catalytic domain. Poly(ADP-ribose) polymerase catalyses the covalent attachment of ADP-ribose units from NAD+ to itself and to a limited number of other DNA binding proteins, which decreases their affinity for DNA. Poly(ADP-ribose) polymerase is a regulatory component induced by DNA damage. The carboxyl-terminal region is the most highly conserved region of the protein. Experiments have shown that a carboxyl 40 kDa fragment is still catalytically active. References: 3 Pubmed Links Taxa: Eukaryota Status: Alignment from source Created: 11-Apr-2003 Representative: Consensus PSSM: 215 columns Aligned: 6 rows Proteins: [Click here for CDART summary of Proteins containing pfam00644] with Cn3D using Virtual Bonds (To display structure, download Cn3D) View 3D Structure width 60 color at 2.0 bits View Alignment All Hypertext of the most diverse members Subset Rows up to 10 20 30 40 5.0 10 1 LKCHLEFYDKDSE----EFSILRQYVKNTHASTHKAYDLK-----IVEVFRVSRQG 47 consensus 136 LRTDIKVVDKDSB----EAKLIKQYVKNTHAATHNAYDLK-----VVEIFRIEREG 182 1EFY A 1353140 304 LPCHLEPVSEEIAgkigDCLAMRGPTHCYKLSLIDAFELKdpneipteaFVEVQEVPKKR 363 gi 1709740 421 LNCGLTPVGNDSE----EFSMVANYMENTHAKTHSGYTVE-----IAQLFRASRAV 467 779 IKTQLVALDKNSE----EFSILSQYVKNTHASTHKSYDLK------IVDVFKVSRQG 825 gi 548585 gi 1709741 782 LETKLEPLDKNSE----EYILLQKYVKNTHAETHKLYDLE-----VVDIFKVARQG 828 80 90 100 70 ...,*....|....*....|.....*....|..... 48 EARRFKPFKKL----HNRRLLWHGSRLTNFAGILSQGLRIAPPEAPVTGYMFGKGVYFAD 103 consensus 183 ESQRYKPFKQL----HNRQLLWHGSRTTNFAGILSQGLRIAPPEAFYTGYMFGKGIYFAD 238 1353140 364 GRKSTKTAAPTVPPPTTKRLLWHGTRVTNVFSILMNGLQF--PVGDRCGLMFGNGVYFAN 421 1709740 468 EADRFQQFSSS----KNRMLLWHGSRLTNWAGILSQGLRIAPFEAPVTGYMFGKGVYFAD 523 gi 548585 826 EARREKPEKKL----HNRKLLWHGSRLTNEVGILSHGLRIAPPEAPPTGYMFGKGIYFAD 881 1709741 829 EARRYKPFKKL----HNRRLLWHGSRLINFAGILSHGLKIRPFEAPVTGYMFGKGIYFAD 884 150 160 170 130 140 104 MVSKSANYCCTSQANSTGLMLLCEYALGD---MYELTIAKY-ITKLPNGKHSVKGRGKTA 159 consensus 239 MVSKSANYCHTSQADPIGLILLGEVALGN---MYELKNASH-ITKLPKGKHSVKGLGKTA 294 gi 1353140 421 VFTKSANYC-CPEASERVFMLLCEVETANPLVLYESEIDAD-EKMEKAKKTSVYAAGKHT 479 1709740 524 KESKSANYCYANTGANDGVLLLCEVALGD---MNELLYSDYNADNLPPGKLSTKGVCKTA 580 gi 548585 882 MVSESANYCCTSQQNSTGLMLLSEVALGD---MMECTSAKY-INKLSNNKHSCFGRGRTM 937 gi 1709741 885 MVSKSANYCCTSHHNSTGLMLLSEVALGD---MMECTAAKY-VTKLPNDKHSCFGRGRTM 940 200 210 220 230*....|....*....|....*....|....*....|....*.... 160 PNPTES-ITL-DGVEVPLGNPIETIELKTSLLYNEYIVYNVEQVKIKYVLRVKFNYKT 215 consensus 295 PDPTAT-TTL-DGVEVPLGNGISTGINDTCLLYNEYIVYDVAQVNLKYLLKLKFNYKT 350 gi 1353140 480 FRDT---VEI-NGIPAFKSN-LETIEEETRLLYDEYVMFNKEHFKIKYVVEVKVDRLT 532 1709740 581 PNPSEA-QTLeDGVVVPLGKPVERSCSKGMLLYNEYIVYNVEQIKMRYVIQVKFNYKH 637 548585 938 PDPTKSyIRS-DGVEIPYGETITDEHLKSSLLYNEYIVYDVAQVNIQYLFRMEFKYSY 994

Help | Disclaimer | Write to the Help Desk

gi 1709741 941 PNFSES-IIR@DGVEIFLGKPITNDSLKSSLLYNEFIIYDIAQVNIQYMLRMNFKYK- 996

Bo

Details

Exhibit K 09/843, 159





SINC	BI	Pu	ID W	16a		of Medicine
PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy
Search PubMed	¥	for				Go Clear
		Limits	Preview/Index	History		Clipboard
About Entrez	7			,		
`	`' I	Display Abstract	<u> </u>	Show: 20 ▼	Sort	Send to

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources Order Documents NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Privacy Policy

1: Proc Natl Acad Sci U S A. 1996 Jul 23;93(15):7481-5.

Related Articles, Links

MIMO

Text

FREE full text article at www.pnas.org in PubMed Central

Structure of the catalytic fragment of poly(AD-ribose) polymerase from chicken.

Ruf A, Mennissier de Murcia J, de Murcia G, Schulz GE.

Institut fur Organische Chemie und Biochemie, Freiburg im Breisgau, Germany.

The crystal structures of the catalytic fragment of chicken poly(ADP-ribose) polymerase [NAD+ADP-ribosyltransferase; NAD+:poly(adenosine-diphosphate-D-ribosyl)-acceptor ADP-D-ribosyltransferase, EC 2.4.2.30] with and without a nicotinamide-analogue inhibitor have been elucidated. Because this enzyme is involved in the regulation of DNA repair, its inhibitors are of interest for cancer therapy. The inhibitor shows the nicotinamide site and also suggests the adenosine site. The enzyme is structurally related to bacterial ADP-ribosylating toxins but contains an additional alpha-helical domain that is suggested to relay the activation signal issued on binding to damaged DNA.

PMID: 8755499 [PubMed - indexed for MEDLINE]

Display Abstract

Show: 20

Sort

Send to Text

■

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer

Jul 8 2003 10:56:01





Clipboard

Send to

63			
163	M	CRI	
	1 J V	டும்.	

PubMed Nucleotide Search PubMed ₹

for

Display

Limits Preview/Index PMC

History

Sort

Taxonomy Go Clear

OMIM Bo

Details

About Entrez

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut

Related Resources Order Documents NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Privacy Policy

1: Gene. 1993 Dec 31;137(2):293-7.

Abstract

Related Articles, Links

Text

Isolation of the poly(ADP-ribose) polymerase-encoding cDNA from Xenopus laevis: phylogenetic conservation of the functional domains.

Uchida K, Uchida M, Hanai S, Ozawa Y, Ami Y, Kushida S, Miwa M.

Show: 20

Department of Biochemistry, University of Tsukuba, Japan.

The complete nucleotide (nt) sequence of the Xenopus laevis poly(ADP-ribose) polymerase (PARP)encoding cDNA was determined. The putative X. laevis PARP protein consists of 1008 amino acids (aa) with a molecular weight of 113 kDa. X. laevis PARP shares 74, 83, 73, 78 and 42% aa sequence homology with the human, bovine, mouse, chicken and Drosophila melanogaster PARPs, respectively. Comparison of the PARP aa sequences among these species showed conservation of two zinc-finger motifs in the DNA-binding domain, and an NAD-binding motif and a Rossmann fold in the catalytic domain. The first Leu of the putative leucine zipper of D. melanogaster PARP is substituted to Lys in X. laevis PARP. All the Glu residues in the leucine zipper are conserved in these six species.

PMID: 8299962 [PubMed - indexed for MEDLINE]


Display **Abstract** Sort Send to

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

> > .bd 8 2003 10-56-01







PubMed Nucleotide Protein Genome PMC Taxonomy OMIM Bo: Search PubMed • Go Clear for Limits Preview/Index Clipboard History Details About Entrez Show: 20 Abstract Display Sort Send to Text Text Version 1: Biochimie. 1995;77(6):456-61. Related Articles, Links Entrez PubMed ELSEVIER Overview Help | FAQ FULLSTEATHERS Tutorial New/Noteworthy Poly(ADP-ribose) polymerase: structure-function relationship. E-Utilities Masson M, Rolli V, Dantzer F, Trucco C, Schreiber V, Fribourg S, Molinete M, Ruf A, Miranda PubMed Services EA, Niedergang C, et al. Journals Database MeSH Database Single Citation Matcher Ecole Superieure de Biotechnologie de Strasbourg, UPR 9003 du CNRS, Illkirch, France. Batch Citation Matcher Clinical Queries Dissection of the human poly(ADP-ribose) polymerase (PARP) molecule in terms of its structure-LinkOut Cubby function relationship has proved to be an essential step towards understanding the biological role of poly(ADP-ribosylation) as a cellular response to DNA damage in eukaryotes. Current approaches Related Resources aimed at elucidating the implication of this multifunctional enzyme in the maintenance of the genomic Order Documents integrity will be presented. **NLM Gateway** TOXNET Consumer Health PMID: 7578429 [PubMed - indexed for MEDLINE] Clinical Alerts
Clinical Trials.gov PubMed Central Privacy Policy Display Abstract Sort Send to **Text**

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

> > Sep 4 2003 10:00:42







SO NCBI		AD M	GU	of Medicin		
	eotide Protein	Genome	Structure	PMC Taxonomy	OMIM	
Search PubMed	for		······································	Go Clear		
	Limits	Preview/Index	History	Clipboar	d Details	
About Entrez				Contraction of the contraction o		
`~ 	Display Abstra	ict 🔻	Show: 20 ▼	Sort <u>▼</u> §Se	nd to Text	
Text Version						
Entrez PubMed	1: Mol Cell Bioch	nem. 1994 Sep;138(1-2):15-24.		Related Articles, Lir	
Overview Help FAQ	Structure o	nd function of poly	(ADD riboso) r	a al remanda de		
Tutorial New/Noteworthy	Structure a	na function of poly	(ADI -House) j	olymei ase.		
E-Utilities		, Schreiber V, Moline	te M, Saulier B,	Poch O, Masson M, N	Niedergang C, Menissie	
PubMed_Services	de Murcia J.					
Journals Database MeSH Database				nite de Cancerogenese		
Single Citation Matcher Batch Citation Matcher	Moleculaire e France.	et Structurale, Centre N	ational de la Rec	herche Scientifique, IIII	kirch-Graffenstaden,	
Clinical Queries LinkOut	Tance.					
Cubby					of systems developed by	
Related Resources				nmental and endogenou ent of genetic and mole		
Order Documents NLM Gateway	elucidate the structure and the function of this abundant and highly conserved enzyme. This article					
TOXNET Consumer Health	summarizes o	our present knowledge i	n this field.			
Clinical Alerts ClinicalTrials.gov	Publication T					
PubMed Central	• Revie	ew ew, Tutorial				
Privacy Policy	• Revie	zw, rutoriai				
	PMID: 78984	58 [PubMed - indexed	for MEDLINE]			

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

Sep 4 2003 10:00:42

Overview

Tutorial New/Noteworthy

E-Utilities

PubMed Services Journals Database

MeSH Database Single Citation Matcher Batch Citation Matcher

Clinical Queries

Related Resources

Order Documents

NLM Gateway

Clinical Alerts

Privacy Policy

Clinical Trials.gov

PubMed Central

TOXNET Consumer Health

LinkOut Cubby

Help | FAQ

8xh.4.60 09/843,159
National





		*		*				
PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	MIMO	В
Search PubMed	∑ i	for			Go	Clear		
		Limits	Preview/Index	History	r	Clipboard		Details
About Entrez	1							
•	~~´ Di	splay Abstract	¥	Show: 20 💌	Sort	Send to	Text	ĮŦ
Text Version				,				

1: J Biol Chem. 1993 Apr 25;268(12):8529-35. Entrez PubMed

Related Articles, Links

FREE full text article at www.jbc.org

> Identification of potential active-site residues in the human poly(ADP-ribose) polymerase.

Simonin F, Poch O, Delarue M, de Murcia G.

Unite propre de recherche de Cancerogenese et de Mutagenese Moleculaire et Structurale, Centre National de la Recherche Scientifique, Strasbourg, France.

The carboxyl-terminal catalytic domain of the human poly(ADP-ribose) polymerase (PARP) exhibits sequence homology with the NAD(P)(+)-dependent leucine and glutamate dehydrogenases. To clarify the role played by some conserved residues between PARP and NAD(P)(+)-dependent dehydrogenases, point mutations were introduced into the whole enzyme context. Non-conservative mutations of Lys-893 (K893I) and Asp-993 (D993A) completely inactivate human PARP, whereas conservative and nonconservative mutations of Asp-914 (D914E and D914A, respectively) and Lys-953 (K953R and K953I, respectively) partially alter PARP activity. The consequences of conservative substitution of Lys-893 and Asp-993 on the kinetic properties of human poly(ADP-ribose) polymerase enzyme and the polymer it synthesizes suggest that these 2 amino acids are directly involved in the covalent attachment of the first ADP-ribosyl residue from NAD+ onto the acceptor amino acid. In addition, the recent resolution of the three-dimensional structure of the NAD(+)-linked glutamate dehydrogenase from Clostridium symbiosum (Baker, P.J., Britton, K.L., Engel, P.C., Farrants, G.W., Lilley, K.S., Rice, D.W., and Stillman, T.J. (1992) Proteins 12, 75-86) strongly supports our alignment with leucine and glutamate dehydrogenases and provides an interesting structural framework for the analysis of our results of site-directed mutagenesis.


PMID: 8473297 [PubMed - indexed for MEDLINE]

Show: 20 Display Abstract Send to

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

> > Sen 4 2003 10:00:42





Send to

Protein PMO Taxonomy Search PubMed V Go Clear for Limits Preview/Index History Clipboard

About Entrez

Text Version

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services Journals Database MeSH Database Single Citation Matcher Batch Citation Matcher Clinical Queries LinkOut Cubby

Related Resources Order Documents NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Privacy Policy

1: Biochemistry. 1997 Oct 7;36(40):12147-54.

Related Articles, Links

Details

OMBA

Text



Abstract

Display

Random mutagenesis of the poly(ADP-ribose) polymerase catalytic domain reveals amino acids involved in polymer branching.

Sort

Show: 20

Rolli V, O'Farrell M, Menissier-de Murcia J, de Murcia G.

Ecole Superieure de Biotechnologie de Strasbourg, UPR A9003 du CNRS, Illkirch-Graffenstaden, France.

Poly(ADP-ribose) polymerase (PARP) is a multifunctional nuclear zinc finger protein which participates in the immediate response of mammalian cells exposed to DNA damaging agents. Given the complexity of the poly(ADP-ribosylation) reaction, we developed a large-scale screening procedure in Escherichia coli to identify randomly amino acids involved in the various aspects of this mechanism. Random mutations were generated by the polymerase chain reaction in a cDNA sequence covering most of the catalytic domain. Out of 26 individual mutations that diversely inactivated the full-length PARP, 22 were found at conserved positions in the primary structure, and 24 were located in the core domain formed by two beta-sheets containing the active site. Most of the PARP mutants were altered in poly(ADP-ribose) elongation and/or branching. The spatial proximity of some residues involved in chain elongation (E988) and branching (Y986) suggests a proximity or a superposition of these two catalytic sites. Other residues affected in branching were located at the surface of the molecule (R847, E923, G972), indicating that protein-protein contacts are necessary for optimal polymer branching. This screening procedure provides a simple and efficient method to explore further the structure-function relationship of the enzyme.

PMID: 9315851 [PubMed - indexed for MEDLINE]

Display Abstract Show: 20 Sort Send to Text

> Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Freedom of Information Act | Disclaimer

> > Jul S 2003 10:56:01